





Review Article

Moderated Starch Nutrition for a Sustainable Dairy Industry: A Refined Pragmatic Wisdom

Akbar Nikkhah^{1*} , and Masoud Alimirzaei² 

¹ Chief Highly Distinguished Professor and Nutritional Scientist, National Elites Foundation, Iran

² Aras-Dam Arsham Co., Poldasht, West Azarbayjan, Iran

* **Corresponding author:** Akbar Nikkhah, Chief Highly Distinguished Professor and Nutritional Scientist, National Elites Foundation, Iran. Email: anikkha@yahoo.com

ARTICLE INFO

Article History:

Received: 03/01/2025

Revised: 29/01/2025

Accepted: 25/02/2025

Published: 24/03/2025



Keywords:

Barley grain

Corn grain

Dairy cow

Health

Longevity

Nutrition

Starch

ABSTRACT

Introduction: Increased cow loss and treatment costs, and reduced longevity and sustainability in the world dairy industry may provide an insight into rethinking the starch feeding in dairy cows. The objective of this mini-review article was to underline the importance of moderated starch nutrition on the sustainable productivity and health of high-producing dairy cows. Increased milk production per cow through both genetic selection and nutritional programs may have led to decreased longevity and higher culling rates in commercial dairy farms. Starch is the primary energy source for lactating dairy cows, supplied mainly by cereal grains such as barley and corn. Feeding higher rates of starch, especially from barley grain with high rumen degradability can lead to health issues such as subacute rumen acidosis, causing economic losses. Reviewing the literature and large dairy herd observations, moderated starch feeding to modern dairy cows without major negative effects on milk yield and cow health should become a new pragmatic wisdom. It is suggested that the short and long-term effects of feeding varying amounts of different starch grains in high-merit dairy cows should be the focus of future research attention.

1. Introduction

Carbohydrates are the main source of energy for lactating dairy cows (about 70% of diet dry matter)¹. Feeding energy-dense diets to support the energy requirements of high-producing dairy cows is a common nutritional strategy in many farms. However, greater starch loads in the rumen may lead to ruminal acidosis and subsequent health issues². Hence, managing and moderating starch nutrition would be a logical practice to optimize dairy cow productivity and sustainability². In addition, limited starch intake has been recommended to increase dry matter intake postpartum^{2,3}. In this review article, a practical perspective was delineated to help optimize dairy profitability and sustainability.

1.1. Starch nutrition dynamics and strategies

Starch is the main component of cereal grains with different inherent features. Cereal grains are different in their starch content and degradability which can impact rumen physiology and animal performance³. Corn and barley grains

are two main sources of starch in dairy diets. Corn grain contains about 70-75% starch and barley grain contains about 50-55%, reflecting their energy density^{3,4}. Starch degradability from corn and barley grains is highly related to the proteinaceous matrix that surrounds starch granules⁴. Given that, corn grain has an extended protein matrix, its degradability is slower than barley grain⁴. Moreover, the processing method (warm vs. cold) has a significant influence on starch availability for the rumen microorganisms and volatile fatty acid (VFA) production⁵. Feeding higher amounts of starchy concentrates is a common practice in many dairy farms to meet the energy requirements of high-producing dairy cows. However, overfeeding starch may predispose cows to metabolic disorders such as rumen acidosis. Decreased dry matter intake (DMI) and milk production and compromised cow welfare via increased prevalence of laminitis are considered as the common consequence of excessive grain feeding in dairy cows^{2,5}. Increased rates of cow loss and

Cite this paper as Akbar Nikkhah, Chief Highly Distinguished Professor and Nutritional Scientist, National Elites Foundation, Iran. Farm Animal Health and Nutrition. 2025; 4(1): 6-8. DOI: 10.58803/fahn.v4i1.66



The Author(s). Published by Rovedar. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

treatment costs, and reduced longevity and sustainability in the world dairy industry may provide an insight into rethinking the starch feeding in dairy cows².

Starch degradability has an important role in nutrient availability in the rumen and post-ruminal starch assimilation. In addition, the optimum level of starch in dairy cattle diets is closely related to dietary neutral detergent fiber (NDF), because both affect DMI, milk yield as well as rumen health⁶. Therefore, optimizing starch nutrition for dairy cows is an art that farm nutritionists should develop and practice. Recently, it was suggested that reduced-starch diets may have benefits for dairy cows⁷. Increased milk yield was reported in cows fed a lower starch diet (22.1%) vs. cows fed a higher starch diet (28.3%) in the fresh period from d 1 to 23 post-calving⁷. In another study, increasing dietary starch (DM basis) by replacing corn grain with beet pulp from 17.7% (low starch) to 24.6% (high starch) did not affect the performance of dairy cows on 118 d in milk⁸. Other studies have shown that feeding a higher-starch diet (26.2% DM basis) relative to a lower-starch diet (21.5% DM basis) in early lactation increased raw milk yield; however, energy-corrected milk yield was not different⁹. It is important to note that cows fed the higher-starch diet lost less body condition score (BCS) than those fed the lower-starch diet^{7,9}. Greater non-esterified fatty acids and β -hydroxybutyrate were reported for low-starch-fed cows, indicating greater body fat mobilization and BCS loss in such cows¹⁰. As such, high-starch-fed cows (27.2 %, DM basis) had reduced systemic inflammation and increased milk production than low-starch-fed cows (25.1 %, DM basis)¹¹. Nonetheless, given that grains comprise up to 30-40% of dairy diets (DM basis), feeding higher starch may be risky, especially when rapidly fermentable cereals such as wheat and barley grains are fed³. Including a maximum of 35% grains (diet DM) even at high levels of production would be watchful³. From a pragmatic point of view, starch inclusion should not exceed 25-28% of diet DM¹².

1.2. Grain type and its relation to grain level

In addition to starch levels, the grain type has a key role in determining starch and NDF concentrations in dairy diets. Furthermore, milk production may be affected by using different grain sources in the diet. It seems that barley grain may provide a more synchronous release of energy and protein than corn grain¹³. It has been demonstrated that microbial nitrogen (N) flow to the duodenum was greater in barley-based diets than in oats-based diet¹⁴. However, such an advantage may disappear when the rumen pH is below the normal range (< 5.8-6.0) because of increased microbial maintenance requirements and wastage¹³. Increased endotoxin levels due to feeding higher levels of highly fermentable carbohydrates can lead to inflammatory responses, immune dysfunction, and reduced longevity. Such conditions may be exacerbated when barley grain is overprocessed¹³. Reduced rumen pH was observed for cows fed flatly rolled barley relative to those fed coarsely rolled barley¹⁵. The authors mentioned that starch digestibility had more correlation with rumen pH than with effective NDF intake. In a recent study investigating the effects of feeding wheat grain or corn grain on dairy cows' performance, a wheat-based diet negatively affected milk production¹⁶. Furthermore, the time period that the rumen pH remained below 6.0 was longer in the wheat-fed cows than in

the corn-fed cows. In line with these results, Oba and Allen (2003) indicated that in corn-based diets, when high starch diets are fed to dairy cows (31%, DM basis), reducing starch fermentation rate by choosing a proper processing method such as dry grinding relative to moisturizing may increase milk yield. A recent large dairy herd study by the current authors demonstrated that decreasing dietary cereal grains from 39% to 34% of diet DM increased milk production (unpublished data; Aras-Dam Arsham, CO., Poldasht, Iran). Interestingly, the original diet had 82% barley grain and 18% corn grain (% of total grain); whereas the refined diet had 38% barley grain and 62% corn grain. In accordance with these farm outcomes, relatively replacing barley grain with corn grain and decreasing total dietary grains by feeding more protein may improve high-merit dairy cow performance.

2. Conclusion

Starch levels and sources in dairy diets play significant roles in determining cow health, productivity, and longevity. It seems that even at high production levels, starch is currently overfed to high-merit dairy cows. The higher starch levels, especially when readily fermentable grains such as barley and wheat are fed, may lead to chronic metabolic disorders such as subacute rumen acidosis, causing serious health and productive issues. It is recommended that starch levels should be controlled in dairy diets, particularly when grain types and processing methods are in favor of higher acid production in the rumen and subsequent health issues. Starch nutrition in high-merit dairy cows demands the development and practice of new pragmatic wisdom. Investigating the short- and long-term effects of feeding varying amounts of different starchy grains in high-merit dairy cows requires future research.

Declarations

Competing interests

There is no competing of interest.

Funding

There is no funding for this study.

Availability of data and materials

Data from the study are available according to a reasonable request.

Authors' contributions

Akbar Nikkhah led the study and conceptualized, strategized, reviewed, edited, and prepared the final manuscript. The initial draft was developed and prepared by Masoud Alimirzaei. The final manuscript was checked by all authors.

Ethical considerations

The authors have made necessary ethical considerations (e.g., plagiarism, consent to publish, misconduct, data fabrication and/or falsification, double publication and/or submission, and redundancy).

Acknowledgments

Profound thanks to Aras-Dam Arsham Co. (Poldasht, West Azarbayjan, Iran).

References

1. Oba M, and Kammes-Main K. Symposium review: Effects of carbohydrate digestion on feed intake and fuel supply. *J Dairy Sci.* 2023; 106(3): 2153-2160. DOI: [10.3168/jds.2022-22420](https://doi.org/10.3168/jds.2022-22420)
2. Nikkhah A. Controlled starch feeding to rescue the dairy industry. *MAVS.* 2016; 1(1): 1-2. Available at: <https://scientiaricerca.com/srmavs/SRMAVS-01-00004.php>
3. Nikkhah A. Optimizing starch utilization: the postmodern art of dairy cow management. *MAVS.* 2017; 1(3): 1-2. Available at: <https://scientiaricerca.com/srmavs/SRMAVS-01-00017.php>
4. Oba M, and Allen MS. Effects of corn grain conservation method on feeding behavior and productivity of lactating dairy cows at two dietary starch concentrations. *J Dairy Sci.* 2003; 86(1): 174-183. DOI: [10.3168/jds.S0022-0302\(03\)73598-X](https://doi.org/10.3168/jds.S0022-0302(03)73598-X)
5. Dehghan-banadaki M, Corbett R, and Oba M. Effects of barley grain processing on productivity of cattle. *Anim Feed Sci Technol.* 2007; 137(1-2): 1-24. DOI: [10.1016/j.anifeedsci.2006.11.021](https://doi.org/10.1016/j.anifeedsci.2006.11.021)
6. National academics of science, engineering, and medicine (NASEM). 2021. Nutrient requirements of dairy cattle. 8th revised ed. National Academic Press.
7. Shi WCE, Knoblock KV, Murphy TC, Bruinje I, Yoon DJ, and Oba M. Effects of supplementing a *Saccharomyces cerevisiae* fermentation product during the periparturient period on performance of dairy cows fed fresh diets differing in starch content. *J Dairy Sci.* 2019; 102(4): 3082-3096. DOI: [10.3168/jds.2018-15307](https://doi.org/10.3168/jds.2018-15307)
8. Dann HM, Tucker HA, Cotanch KW, Mooney CS, Grant RJ, Eguchi T. Evaluation of low starch diets for lactating Holstein dairy cows. *J Dairy Sci.* 2014; 97(11): 7151-DOI: [10.3168/jds.2014-8341](https://doi.org/10.3168/jds.2014-8341)
9. McCarthy MM, Yosui T, Ryan CM, Mechor GD, and Overton TR. Performance of early-lactation dairy cows as affected by dietary starch and monensin supplementation. *J Dairy Sci* 2015; 98(5): 3335-3350. DOI: [10.3168/jds.2014-8820](https://doi.org/10.3168/jds.2014-8820)
10. Piccioli-Cappelli F, Seal CJ, Parker DS, Looor JJ, Minuti A, Lopreiato V, et al. *J Dairy Sci.* 2022; 105(2): 1131-1149. DOI: [10.3168/jds.2021-20539](https://doi.org/10.3168/jds.2021-20539)
11. Haisan J, Inabo Y, and Oba M. Effects of pre-and postpartum dietary starch content on productivity, plasma energy metabolites, and serum inflammation indicators of dairy cows. *J Dairy Sci.* 2020; 104(4): 4362-4374. DOI: [10.3168/jds.2020-19611](https://doi.org/10.3168/jds.2020-19611)
12. Nikkhah A. Optimizing starch nutrition for rhythmic dairy cattle: The sustaining economic and environmental challenges of today's industry. *Iran J Applied Anim Sci.* 2021; 11(2): 207-215. Available at: https://journals.iau.ir/article_682225_675c0b028da41a33b062ef159d13ca94.pdf
13. Nikkhah A. Barley grain for ruminants: A global treasure or tragedy. *J Anim Sci Biotechnol.* 2012; 3: 22. DOI: [10.1186/2049-1891-3-22](https://doi.org/10.1186/2049-1891-3-22)
14. Gozho GN, and Mutsvangwa T. Influence of carbohydrate source on ruminal fermentation characteristics, performance, and microbial protein synthesis in dairy cows. *J Dairy Sci.* 2008; 91(7): 2726-2735. DOI: [10.3168/jds.2007-0809](https://doi.org/10.3168/jds.2007-0809)
15. Yang WZ, Beauchemin KA, and Rde LM. Effects of grain processing, forage to concentrate ratio, and forage particle size on rumen pH and digestion by dairy cows. *J Dairy Sci.* 2001; 84(10): 2203-2216. DOI: [10.3168/jds.S0022-0302\(01\)74667-X](https://doi.org/10.3168/jds.S0022-0302(01)74667-X)
16. Moate OJ, Williams MH, Deighton MC, Hannah BE, Ribaux G, Morris L, et al. Effects of feeding wheat or corn and of rumen fistulation on milk production and methane emissions of dairy cows. *Anim Prod Sci.* 2019; 59(5): 891-905. DOI: [10.1071/AN17433](https://doi.org/10.1071/AN17433)